

**Supplementary Table 3.** Compliance with ERAS interventions

ERAS intervention	Agreement rate (%) <sup>a</sup>	Data not available (n)
Preadmission counselling	100	1
Carbohydrate loading	97.1	1
Immunonutrition	0	11
Surgical access		0
Open	11.7	
Laparoscopic	63.2	
Laparoscopic converted to open	11.1	
Hybrid	14.0	
Bowel preparation	91.2	0
Oral antibiotics	1.2	0
Liquids up to 3 hr prior to surgery	73.7	0
Epidural	5.3	0
Multimodal pain management	88.2	1
NSAIDs	31.8	1
Preoperative iron transfusion for anemia	6.3	12
Postoperative antiemetics	85.4	0
Mobilization POD 0–1	83.0	0
Mobilization BD POD 0–1	82.9	19
IV fluids discontinued POD 0–1	40.1	4
Upgraded to solid diet POD 0–2	71.4	0
IDC		
IDC out POD 0–2	79.5	0
Failed TOV, IDC reinsertion required	8.8	0
NGT		
NGT out POD 0–2	3.5	
NGT out POD ≥ 3	8.8	
No NGT	87.7	
NGT reinsertion	14.6	0
Drain	1.0	
No drain	54.7	
Drain removed POD 0–2	10.6	
Drain removed POD 3–4	17.6	
Drain removed POD ≥ 5	14.6	0

ERAS, Enhanced Recovery After Surgery; NSAIDs, nonsteroidal anti-inflammatory drugs; POD, postoperative day; BD, twice a day; IV, intravenous; IDC, indwelling catheter; TOV, trial of void; NGT, nasogastric tube.

<sup>a</sup>Does not include missing data.